

Hakai Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_079090.2

Catalog No. A302-968A-T GenelD 79872

Lot No. A302-968A-T-1

APPLICATIONS	IP, IHC
SPECIES REACTIVITY	Human
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to Hakai immobilized on solid support.

The epitope recognized by A302-968A-T maps to a region between residue 1 and 50 of human E3 ubiquitin-protein ligase Hakai using the numbering given in entry NP_079090.2 (GeneID 79872).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended



Immunoprecipitation 2 – 5 µg/mg lysate

Immunohistochemistry 1:500 – 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.

ADDITIONAL INFO <https://www.bethyl.com/product/A302-968A-T>

Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human Hakai by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-Hakai antibody A302-968A used for IP at 3 µg/mg lysate. Hakai was also immunoprecipitated by rabbit anti-Hakai antibody A302-969A, which recognizes a downstream epitope. For blotting immunoprecipitated Hakai, A302-969A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.

Detection of human Hakai by immunohistochemistry. *Sample:* FFPE section of human breast carcinoma. *Antibody:* Affinity purified rabbit anti-Hakai (Cat. No. A302-968A Lot1) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* DAB