RLF Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_036553.2

Catalog No. A303-207A-M GeneID 6018

Lot No. A303-207A-M-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human

AMOUNT 100 μl (10 blots)

CONCENTRATION 100 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to RLF immobilized on solid support.

The epitope recognized by A303-207A-M maps to a region between residue 500 and 550 of human Rearranged L-Myc Fusion using the numbering given in entry NP_036553.2 (GenelD

6018).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation The antibody contained within A303-207A-M has been qualified for

use in immunoprecipitation; however, we recommend using the alternative formulation of this antibody found as product A303-207A.

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A303-207A-M

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



RLF Antibody A303-207A-M



Detection of human RLF by western blot. Samples: Whole cell lysate from HeLa (15 and 50 μg) and HEK293T (T; 50 μg) cells. Antibody: Affinity purified rabbit anti-RLF antibody A303-207A-M used at 1:1000. Detection: Chemiluminescence with an exposure time of 3 minutes.